

Periodic Oscillation for Cohen-Grossberg-type Neural Networks with Neutral Time-varying Delays

Chuanzhi Bai*

Abstract: In this paper, a Cohen-Grossberg-type model of neural networks with neutral time-varying delays is investigated by using the continuation theorem of Mawhin's coincidence degree theory and some analysis techniques. Without assuming the continuous differentiability of time-varying delays, sufficient conditions for the existence of the periodic solutions are given. The result of this paper is new and extent previously known result.

Keywords: Cohen-Grossberg type neural networks; Neutral delay; Coincidence degree theory; Periodic solution

1. INTRODUCTION

In recent years, the cellular neural networks have been extensively studied and applied in many different fields such as signal and image processing, pattern recognition and optimization. In implementation of networks, time delays are inevitably encountered because of the finite switching speed of amplifiers. Thus, it is very important to investigate the dynamics of delay neural networks. In theory and application, the existence of periodic solutions of neural networks model is of great interest [1-9]. Very recently, Gui, Ge and Yang [10] investigated the existence of periodic solutions of Hopfield networks model with neutral delay by means of an abstract continuous theorem of k -set contractive operator and some analysis techniques. But they do require the time-varying delays $\tau_{ij}(t)$ and $\sigma_{ij}(t)$ are continuously differentiable. Furthermore, the criterion for the existence of periodic solutions of Hopfield neural networks model in [10] depends on the $\dot{\tau}_{ij}$ and $\dot{\sigma}_{ij}$.

In this paper, we consider the following Cohen-Grossberg type neural networks with neutral time-varying delays:

$$\dot{x}_i(t) = -a_i(x_i(t)) \left[b_i(x_i(t)) - \sum_{j=1}^n a_{ij}(t) f_j(x_j(t - \tau_{ij}(t))) - \sum_{j=1}^n b_{ij}(t) g_j(\dot{x}_j(t - \sigma_{ij}(t))) - J_i(t) \right], \quad (1)$$

where $i = 1, 2, \dots, n$, $x_i(t)$ denotes the potential (or voltage) of cell i at time t ; $a_i(x_i(t))$ represent an amplification function; $b_i(x_i(t))$ be include a constant term indicating a fixed input the network; $a_{ij}(t)$ and $b_{ij}(t)$ represent the delayed strengths of connectivity and neutral delayed strengths of connectivity between cell i and j at time t , respectively; f_j and g_j are the activation functions in system (1); $J_i(t)$ is an external input on the i th unit at time t , in which $J_i : \mathbb{R} \rightarrow \mathbb{R}$, $i = 1, \dots, n$, are continuous periodic functions with period ω ; $\tau_{ij}(t)$, $\sigma_{ij}(t) \geq 0$ correspond to the transmission delays.

Obviously, system (1) is a generalization of the following model for an Hopfield neural networks system with neutral delays in [10]:

$$\dot{x}_i(t) = -b_i x_i + \sum_{j=1}^n a_{ij}(t) g_j(x_j(t - \tau_{ij}(t))) + \sum_{j=1}^n b_{ij}(t) g_j(\dot{x}_j(t - \sigma_{ij}(t))) + J_i(t). \quad (2)$$

* Department of Mathematics, Huaiyin Teachers College, Huaian, Jiangsu 223300, P R China, E-mail: czbai8@sohu.com

In this paper, by using the continuation theorem of coincidence degree theory and some analysis technique, we obtain some new sufficient conditions for the existence of the periodic solutions of system (1). The conditions imposed on the time-varying delays $\tau_{ij}(t)$ and $\sigma_{ij}(t)$ do not need the assumptions of continuously differentiable. Our works in this paper are new and extend previous result in [10].

2. PRELIMINARIES

In this section, we state some notations, definitions and some Lemmas.

Let $A = (a_{ij})_{n \times n}$ be a real $n \times n$ matrix, $A > 0$ ($A \geq 0$) denotes each element a_{ij} is positive (nonnegative, respectively). Let $x = (x_1, x_2, \dots, x_n)^T \in \mathbb{R}^n$ be a vector, $x > 0$ ($x \geq 0$) denotes each element x_i is positive (nonnegative, respectively). For matrices or vectors A and B , $A \geq B$ ($A > B$) means that $A - B \geq 0$ ($A - B > 0$).

Definition 2.1 [11]: Matrix $A = (a_{ij})_{n \times n}$ is said to be a nonsingular M -matrix, if

- (i) $a_{ii} > 0, i = 1, 2, \dots, n$;
- (ii) $a_{ij} \leq 0$, for $i \neq j, i, j = 1, 2, \dots, n$;
- (iii) $A^{-1} \geq 0$.

Let X and Y be normed vector spaces, Let $L : \text{dom}L \subset X \rightarrow Y$ be a linear mapping, L will be called a Fredholm mapping of index zero if $\dim \text{Ker}L = \text{codim} \text{Im}L < +\infty$ and $\text{Im}L$ is closed in Y . If L is a Fredholm mapping of index zero, there exist continuous projectors $P : X \rightarrow X$ and $Q : Y \rightarrow Y$ such that $\text{Im}P = \text{Ker}L$, $\text{Ker}Q = \text{Im}L = \text{Im}(I - Q)$. It follows that mapping $L|_{\text{dom}L \cap \text{Ker}P} : (I - P)X \rightarrow \text{Im}L$ is invertible. We denote the inverse of the mapping by K_p . If Ω is an open bounded subset of X , the mapping N will be called L -compact on $\bar{\Omega}$ if $QN(\bar{\Omega})$ is bounded and $K_p(I - Q)N : \bar{\Omega} \rightarrow X$ is compact. Since $\text{Im}Q$ is isomorphic to $\text{Ker}L$, there exists an isomorphism $J : \text{Im}Q \rightarrow \text{Ker}L$.

Now, we introduce Mawhin's continuation theorem [12, p.40] as follows.

Lemma 2.1: Let X and Y be two Banach spaces, $L : \text{dom}L \rightarrow Y$ be a Fredholm operator with index zero. Assume that Ω is an open bounded set in X , and N is L -compact on $\bar{\Omega}$. If all the following conditions hold:

- (a) for each $\lambda \in (0, 1)$, $x \in \partial \Omega \cap \text{Dom}L$, $Lx \neq \lambda Nx$;
- (b) $QNx \neq 0$ for each $x \in \partial \Omega \cap \text{Ker}L$, and $\deg(JNQ, \Omega \cap \text{Ker}L, 0) \neq 0$,

where J is an isomorphism $J : \text{Im}Q \rightarrow \text{Ker}L$. Then equation $Lx = Nx$ has at least one solution in $\bar{\Omega} \cap \text{Dom}L$.

The following lemmas will be useful to prove our main result in Section 3.

Lemma 2.2 [11]: Assume that A is a nonsingular M -matrix and $Aw \leq d$, then $w \leq A^{-1}d$.

Lemma 2.3 [13]: Let $A = (a_{ij})$ with $a_{ij} \leq 0, i, j = 1, 2, \dots, n, i \neq j$. Then the following statements are equivalent.

- (1) A is an M -matrix.
- (2) There exists a row vector $\eta = (\eta_1, \eta_2, \dots, \eta_n) > (0, 0, \dots, 0)$ such that $\eta A > 0$.
- (3) There exists a column vector $\xi = (\xi_1, \xi_2, \dots, \xi_n)^T > (0, 0, \dots, 0)^T$ such that $A\xi > 0$.

Throughout this paper, we assume that

(H_1) $a_{ij}, b_{ij}, J_j \in C(\mathbb{R}, \mathbb{R}), \tau_{ij}, \sigma_{ij} \in C(\mathbb{R}, \mathbb{R}^+)$ ($\mathbb{R}^+ = [0, \infty)$) are periodic functions with a common period $\omega (> 0)$, $i, j = 1, \dots, n$.

(H₂) $a_i \in C(\mathbb{R}, \mathbb{R})$. Furthermore, there exist positive constants a_{i*} and a_i^* such that

$$a_{i*} \leq a_i(u) \leq a_i^*, \quad \forall u \in \mathbb{R}, \quad i = 1, \dots, n.$$

(H₃) $b_i \in C(\mathbb{R}, \mathbb{R})$. Moreover, there exist positive constants b_{i*} and b_i^* such that

$$b_{i*}u^2 \leq b_i(u) \leq b_i^*u^2, \quad \forall u \in \mathbb{R}, \quad i = 1, \dots, n.$$

(H₄) $f_j, g_j \in C(\mathbb{R}, \mathbb{R})$ are Lipschitzian with Lipschitz constants L_j and l_j respectively, i.e.,

$$|f_j(x) - f_j(y)| \leq L_j|x - y|, \quad |g_j(x) - g_j(y)| \leq l_j|x - y|, \quad \forall x, y \in \mathbb{R}, \quad j = 1, \dots, n.$$

3. EXISTENCE OF PERIODIC SOLUTION

In this section, we will use the continuation theorem of coincidence degree theory to obtain the existence of an ω -periodic solution to system (1).

For convenience, we introduce the following notations:

$$a_{ij}^+ := \max_{t \in [0, \omega]} |a_{ij}(t)|, \quad b_{ij}^+ := \max_{t \in [0, \omega]} |b_{ij}(t)|, \quad J_i^+ := \max_{t \in [0, \omega]} |J_i(t)|, \quad i, j = 1, 2, \dots, n.$$

Theorem 3.1: Let (H₁)-(H₄) hold. Suppose that C and $A - B(C^{-1}D)$ are two nonsingular M -matrix, then system (1) has at least one ω -periodic solution, where

$$\begin{aligned} A &= (\bar{a}_{ij})_{n \times n}, \quad \bar{a}_{ij} = b_{i*} \delta_{ij} - a_{ij}^+ L, \quad B = (\bar{b}_{ij})_{n \times n}, \quad \bar{b}_{ij} = b_{ij}^+ l_j, \\ C &= (\bar{c}_{ij})_{n \times n}, \quad \bar{c}_{ij} = \delta_{ij} - a_i^* b_{ij}^+ l_j, \quad D = (\bar{d}_{ij})_{n \times n}, \quad \bar{d}_{ij} = a_i^* (b_i^* \delta_{ij} + a_{ij}^+ L_j), \quad \delta_{ij} = \begin{cases} 1, & \text{for } i = j, \\ 0, & \text{for } i \neq j. \end{cases} \end{aligned}$$

Proof: Take

$$C_\omega = \{x(t) = (x_1(t), \dots, x_n(t))^T \in C(\mathbb{R}, \mathbb{R}^n) : x_i(t + \omega) \equiv x_i(t), \quad i = 1, \dots, n\},$$

$$C_\omega^1 = \{x(t) = (x_1(t), \dots, x_n(t)) \in C^1(\mathbb{R}, \mathbb{R}^n) : x_i(t + \omega) \equiv x_i(t), \quad i = 1, \dots, n\}.$$

Then C_ω is a Banach space with the norm $\|x\|_0 = \max_{1 \leq i \leq n} \{|x_i|_0\}$, $|x_i|_0 = \max_{t \in [0, \omega]} |x_i(t)|$, C_ω^1 is also a Banach space with the norm $\|x\| = \max\{\|x\|_0, \|x'\|_0\}$.

For each $x = (x_1, \dots, x_n) \in C_\omega^1$, $L : C_\omega^1 \rightarrow C_\omega$ and $N : C_\omega^1 \rightarrow C_\omega$ defined by

$$(Lx)(t) = \frac{dx}{dt} = (\dot{x}_1(t), \dots, \dot{x}_n(t))^T, \quad \text{and}$$

$$(Nx)(t) = \begin{bmatrix} -a_1(x_1(t)) \left(b_1(x_1(t)) - \sum_{j=1}^n a_{1j}(t) f_j(x_j(t - \tau_{1j}(t))) - \sum_{j=1}^n b_{1j}(t) g_j(\dot{x}_j(t - \sigma_{1j}(t))) - J_1(t) \right) \\ \vdots \\ -a_n(x_n(t)) \left(b_n(x_n(t)) - \sum_{j=1}^n a_{nj}(t) f_j(x_j(t - \tau_{nj}(t))) - \sum_{j=1}^n b_{nj}(t) g_j(\dot{x}_j(t - \sigma_{nj}(t))) - J_n(t) \right) \end{bmatrix}.$$

It is easy to see that $\text{Ker}L = \mathbb{R}^n$, $\text{Im}L = \{(x_1(t), \dots, x_n(t))^T \in C_\omega : \int_0^\omega x_i(s) ds = 0, \quad i = 1, \dots, n\}$ is closed in C_ω , and L is a Fredholm mapping of index zero. Define two projectors

$$Px = \frac{1}{\omega} \int_0^\omega x(t) dt = \left(\frac{1}{\omega} \int_0^\omega x_1(t) dt, \dots, \frac{1}{\omega} \int_0^\omega x_n(t) dt \right)^T, \quad x(t) = (x_1(t), \dots, x_n(t))^T \in C_\omega^1,$$

$$Qy = \frac{1}{\omega} \int_0^\omega y(t) dt = \left(\frac{1}{\omega} \int_0^\omega y_1(t) dt, \dots, \frac{1}{\omega} \int_0^\omega y_n(t) dt \right)^T, y(t) = (y_1(t), \dots, y_n(t))^T \in C_\omega.$$

Obviously, P, Q are continuous and satisfy

$$\text{Im}P = \text{Ker}L, \text{Ker}Q = \text{Im}L.$$

Similar to [4], we can define the generalized inverse $K_p : \text{Im}L \rightarrow \text{Ker}P \cap \text{dom}L$ of L and show that N is L -compact on $\bar{\Omega}$ for any open bounded set $\Omega \subset C_\omega^1$.

Now we are in a position to show that there exists an appropriate open, bounded subset Ω , which satisfies all the requirements given in the continuation theorem. According to the operator equation $Lx = \lambda Nx$, $\lambda \in (0, 1)$, we have

$$\dot{x}_i(t) = -\lambda a_i(x_i(t)) \left[b_i(x_i(t)) - \sum_{j=1}^n a_{ij}(t) f_j(x_j(t - \tau_{ij}(t))) - \sum_{j=1}^n b_{ij}(t) g_j(\dot{x}_j(t - \sigma_{ij}(t))) - J_i(t) \right], \quad (3)$$

where $i = 1, \dots, n$. Suppose that $x(t) = (x_1(t), \dots, x_n(t))^T \in C_\omega^1$ is a solution of system (3) for some $\lambda \in (0, 1)$. Hence, there exist $\xi_i \in [0, \omega]$ ($i = 1, \dots, n$) such that $|x_i(\xi_i)| = \max_{t \in [0, \omega]} |x_i(t)| = |x_i|_0$. Thus, $x'_i(\xi_i) = 0$ for $i = 1, \dots, n$. By (3), we have

$$b_i(x_i(\xi_i)) = \sum_{j=1}^n a_{ij}(\xi_i) f_j(x_j(\xi_i - \tau_{ij}(\xi_i))) + \sum_{j=1}^n b_{ij}(\xi_i) g_j(\dot{x}_j(\xi_i - \sigma_{ij}(\xi_i))) + J_i(\xi_i), \quad i = 1, \dots, n. \quad (4)$$

In view of (H_3) , (H_4) and (4), we have

$$\begin{aligned} b_{i*} |x_i|_0 &= b_{i*} |x_i(\xi_i)| \leq |b_i(x_i(\xi_i))| \leq \sum_{j=1}^n |a_{ij}(\xi_i)| |f_j(x_j(\xi_i - \tau_{ij}(\xi_i)))| \\ &\quad + \sum_{j=1}^n |b_{ij}(\xi_i)| |g_j(\dot{x}_j(\xi_i - \sigma_{ij}(\xi_i)))| + |J_i(\xi_i)| \\ &\leq \sum_{j=1}^n a_{ij}^+(L_j |x_j(\xi_i - \tau_{ij}(\xi_i))| + |f_j(0)|) + \sum_{j=1}^n b_{ij}^+(l_j |\dot{x}_j(\xi_i - \sigma_{ij}(\xi_i))| + |g_j(0)|) + J_i^+ \\ &\leq \sum_{j=1}^n (a_{ij}^+ L_j |x_j|_0 + b_{ij}^+ l_j |\dot{x}_j|_0) + \sum_{j=1}^n (a_{ij}^+ |f_j(0)| + b_{ij}^+ |g_j(0)|) + J_i^+, \quad i = 1, \dots, n, \end{aligned}$$

which implies that

$$\sum_{j=1}^n (b_{i*} \delta_{ij} - a_{ij}^+ L_j) |x_j|_0 \leq \sum_{j=1}^n b_{ij}^+ l_j |\dot{x}_j|_0 + \sum_{j=1}^n (a_{ij}^+ |f_j(0)| + b_{ij}^+ |g_j(0)|) + J_i^+, \quad i = 1, \dots, n. \quad (5)$$

The formulas (5) may be rewritten in the form

$$AX \leq BY + h, \quad (6)$$

where $X = (|x_1|_0, \dots, |x_n|_0)^T$, $Y = (|\dot{x}_1|_0, \dots, |\dot{x}_n|_0)^T$, $h = (h_i)_{n \times 1}$, $h_i = \sum_{j=1}^n (a_{ij}^+ |f_j(0)| + b_{ij}^+ |g_j(0)|) + J_i^+$.

Let $\eta_i \in [0, \omega]$ ($i = 1, \dots, n$) such that $|\dot{x}_i(\eta_i)| = \max_{t \in [0, \omega]} |\dot{x}_i(t)| = |\dot{x}_i|_0$. From (3), (H₂), (H₃) and (H₄), we get

$$\begin{aligned} |\dot{x}_i|_0 &= |\dot{x}_i(\eta_i)| \leq a_i(x_i(\eta_i)) \left[|b_i(x_i(\eta_i))| + \sum_{j=1}^n |a_{ij}(\eta_i)| |f_j(x_j(\eta_i - \tau_{ij}(\eta_i)))| \right. \\ &\quad \left. + \sum_{j=1}^n |b_{ij}(\eta_i)| |g_j(\dot{x}_j(\eta_i - \sigma_{ij}(\eta_i)))| + |J_i(\eta_i)| \right] \\ &\leq a_i^* \left[b_i^* |x_i(\eta_i)| + \sum_{j=1}^n a_{ij}^+ (L_j |x_j(\eta_i - \tau_{ij}(\eta_i))| + |f_j(0)|) + \sum_{j=1}^n b_{ij}^+ (l_j |\dot{x}_j(\eta_i - \sigma_{ij}(\eta_i))| + |g_j(0)|) + J_i^+ \right] \\ &\leq a_i^* \left[b_i^* |x_i|_0 + \sum_{j=1}^n (a_{ij}^+ L_j |x_j|_0 + b_{ij}^+ l_j |\dot{x}_j|_0) + \sum_{j=1}^n (a_{ij}^+ |f_j(0)| + b_{ij}^+ |g_j(0)|) + J_i^+ \right], \quad i = 1, \dots, n, \end{aligned}$$

that is

$$\sum_{j=1}^n (\delta_{ij} - a_i^* b_{ij}^+ l_j) |\dot{x}_j|_0 \leq a_i^* \sum_{j=1}^n (b_i^* \delta_{ij} + a_{ij}^+ L_j) |x_j|_0 + a_i^* \sum_{j=1}^n (a_{ij}^+ |f_j(0)| + b_{ij}^+ |g_j(0)|) + a_i^* J_i^+, \quad (7)$$

where $i = 1, \dots, n$. It is easy to know that (7) may be rewritten as

$$CY \leq DX + k. \quad (8)$$

where $k = (k_i)_{n \times 1}$, $k_i = a_i^* \sum_{j=1}^n (a_{ij}^+ |f_j(0)| + b_{ij}^+ |g_j(0)|) + a_i^* J_i^+$. Since C is a nonsingular M -matrix, we have by (8) and Lemma 2.2 that

$$Y \leq C^{-1}DX + C^{-1}k. \quad (9)$$

Substituting (9) into (6), we get

$$(A - B(C^{-1}D))X \leq BC^{-1}k + h := w = (w_1, w_2, \dots, w_n)^T. \quad (10)$$

Since $A - B(C^{-1}D)$ is a nonsingular M -matrix, we have by (10) and Lemma 2.2 that

$$X \leq (A - B(C^{-1}D))^{-1}w := (R_1, R_2, \dots, R_n)^T. \quad (11)$$

Substituting (11) into (9), we obtain

$$Y \leq C^{-1}D(R_1, \dots, R_n)^T + C^{-1}k := (r_1, r_2, \dots, r_n)^T. \quad (12)$$

Since $A - B(C^{-1}D)$ is an M -matrix, we have from Lemma 2.3 that there exists a vector $\varsigma = (\varsigma_1, \varsigma_2, \dots, \varsigma_n)^T > (0, 0, \dots, 0)^T$ such that

$$(A - B(C^{-1}D))\zeta > (0, 0, \dots, 0)^T,$$

which implies that we can choose a constant $p > 1$ such that

$$p(A - B(C^{-1}D))\zeta > w, \text{ and } p\zeta_i > R_i, i = 1, 2, \dots, n. \quad (13)$$

Combine (11) with (8), we get

$$CY \leq D(R_1, \dots, R_n)^T + k := v = (v_1, v_2, \dots, v_n)^T. \quad (14)$$

Noticing that C is an M -matrix, we have from Lemma 2.3 that there exists a vector $\zeta = (\zeta_1, \dots, \zeta_n)^T > (0, 0, \dots, 0)^T$ such that

$$C\zeta > (0, 0, \dots, 0)^T,$$

which implies that we can choose a constant $q > 1$ such that

$$qC\zeta > v, \text{ and } q\zeta_i > r_i, i = 1, 2, \dots, n. \quad (15)$$

Set

$$\bar{\zeta} = (\bar{\zeta}_1, \bar{\zeta}_2, \dots, \bar{\zeta}_n)^T := p\zeta, \quad \bar{\zeta} = (\bar{\zeta}_1, \bar{\zeta}_2, \dots, \bar{\zeta}_n)^T := q\zeta.$$

Then, by (13) and (15), we have

$$\bar{\zeta}_i > R_i, (A - B(C^{-1}D))\bar{\zeta} > w, \bar{\zeta}_i > r_i, \text{ and } C\bar{\zeta} > v, i = 1, 2, \dots, n. \quad (16)$$

Now we take

$$\Omega = \{x(t) = (x_1(t), x_2(t), \dots, x_n(t))^T \in C_\omega^1 : |x_i|_0 < \bar{\zeta}_i, |\dot{x}_i|_0 < \bar{\zeta}_i, i = 1, 2, \dots, n\}.$$

Obviously, the condition (a) of Lemma 2.1 is satisfied. If $x \in \partial \Omega \cap \text{Ker}L = \partial \Omega \cap \mathbb{R}^n$, then $x(t)$ is a constant vector in \mathbb{R}^n , and there exists some $i \in \{1, 2, \dots, n\}$ such that $|x_i| = \bar{\zeta}_i$. It follows that

$$(QNx)_i = -\frac{a_i(x_i)}{\omega} \int_0^\omega \left[b_i(x_i) - \sum_{j=1}^n a_{ij}(t)f_j(x_j) - \sum_{j=1}^n b_{ij}(t)g_j(0) - J_i(t) \right] dt. \quad (17)$$

We claim that

$$|(QNx)_i| > 0. \quad (18)$$

In fact, if $|(QNx)_i| = 0$, i.e.,

$$\int_0^\omega \left[b_i(x_i) - \sum_{j=1}^n f_j(x_j)a_{ij}(t) - \sum_{j=1}^n g_j(0)b_{ij}(t) - J_i(t) \right] dt = 0.$$

Then, there exists some $t_* \in [0, \omega]$ such that

$$b_i(x_i) - \sum_{j=1}^n f_j(x_j)a_{ij}(t_*) - \sum_{j=1}^n g_j(0)b_{ij}(t_*) - J_i(t_*) = 0,$$

which implies that

$$b_{i*}\bar{\zeta}_i = b_{i*}|x_i| \leq |b_i(x_i)| \leq \sum_{j=1}^n |f_j(x_j)|a_{ij}^+ + \sum_{j=1}^n |g_j(0)|b_{ij}^+ + J_i^+$$

$$\begin{aligned} &\leq \sum_{j=1}^n \left(|f_j(x_j) - f_j(0)| + |f_j(0)| \right) a_{ij}^+ + \sum_{j=1}^n |g_j(0)| b_{ij}^+ + J_i^+ \\ &\leq \sum_{j=1}^n a_{ij}^+ L_j \bar{\zeta}_j + \sum_{j=1}^n \left(a_{ij}^+ |f_j(0)| + b_{ij}^+ |g_j(0)| \right) + J_i^+. \end{aligned}$$

This means that

$$(A \bar{\zeta})_i \leq h_i. \quad (19)$$

It is easy to know that $D \bar{\zeta} = (0, 0, \dots, 0)^T$. Since C is a nonsingular M -matrix, we have from $C^{-1} \geq 0$ (Lemma 2.2) that $C^{-1}D \bar{\zeta} \geq (0, 0, \dots, 0)^T$. Thus, we obtain

$$B(C^{-1}D) \bar{\zeta} \geq (0, \dots, 0)^T. \quad (20)$$

Similarly, we have

$$BC^{-1}k \geq (0, \dots, 0)^T. \quad (21)$$

From (19), (20) and (21), we get

$$(A \bar{\zeta})_i \leq h_i + (BC^{-1}k)_i + (B(C^{-1}D) \bar{\zeta})_i = w_i + (B(C^{-1}D) \bar{\zeta})_i.$$

This implies that

$$(A - B(C^{-1}D) \bar{\zeta})_i \leq w_i,$$

which contradicts (16). Hence, (18) holds. Consequently, $QNx, \neq 0$ for each $x \in \partial\Omega \cap \text{Ker}L$.

Furthermore, let $\psi(x, \mu) = \mu(-x) + (1-\mu)JQNx$ ($\mu \in [0, 1]$), then for any $x = (x_1, x_2, \dots, x_n)^T \in \partial\Omega \cap \text{Ker}L$, $(x_1, x_2, \dots, x_n)^T$ is a constant vector in \mathbb{R}^n with $|x_i| = \bar{\zeta}_i$ for some $i \in \{1, \dots, n\}$. It follows that

$$(\Psi(x, \mu))_i = -\mu x_i - (1-\mu) \frac{1}{\omega} a_i(x_i) \int_0^\omega \left[b_i(x_i) - \sum_{j=1}^n f_j(x_j) a_{ij}(t) - \sum_{j=1}^n g_j(0) b_{ij}(t) - J_i(t) \right] dt. \quad (22)$$

We claim that

$$|(\Psi(x, \mu))_i| > 0. \quad (23)$$

If this is not true, then $|(\Psi(x, \mu))_i| = 0$, i.e.,

$$\mu x_i + (1-\mu) \frac{a_i(x_i)}{\omega} \int_0^\omega \left[b_i(x_i) - \sum_{j=1}^n f_j(x_j) a_{ij}(t) - \sum_{j=1}^n g_j(0) b_{ij}(t) - J_i(t) \right] dt = 0.$$

Therefore, there exists some $t^* \in [0, \omega]$ such that

$$\mu x_i + (1-\mu) \frac{a_i(x_i)}{\omega} \left[b_i(x_i) - \sum_{j=1}^n f_j(x_j) a_{ij}(t^*) - \sum_{j=1}^n g_j(0) b_{ij}(t^*) - J_i(t^*) \right] = 0, \quad (24)$$

which implies that

$$x_i \left[b_i(x_i) - \sum_{j=1}^n f_j(x_j) a_{ij}(t^*) - \sum_{j=1}^n g_j(0) b_{ij}(t^*) - J_i(t^*) \right] \leq 0.$$

Thus, we get

$$\begin{aligned} b_{i^*} |x_i|^2 &\leq x_i b_i(x_i) \leq x_i \left[\sum_{j=1}^n f_j(x_j) a_{ij}(t^*) + \sum_{j=1}^n g_j(0) b_{ij}(t^*) + J_i(t^*) \right] \\ &\leq |x_i| \left[\sum_{j=1}^n (|f_j(x_j) - f_j(0)| + |f_j(0)|) a_{ij}^+ + \sum_{j=1}^n g_j(0) b_{ij}^+ + J_i^+ \right] \\ &\leq |x_i| \left[\sum_{j=1}^n a_{ij}^+ L_j \bar{\zeta}_j + \sum_{j=1}^n (a_{ij}^+ |f_j(0)| + b_{ij}^+ |g_j(0)|) + J_i^+ \right], \end{aligned}$$

this means that $(A\bar{\zeta})_i \leq h_i$. By (20) and (21) we obtain

$$((A - BC^{-1}D)\bar{\zeta})_i \leq w_i.$$

which contradicts (16). Hence, (23) holds. By the homotopy invariance theorem, we get

$$\deg\{JQN, \Omega \cap \text{Ker}L, 0\} = \deg\{-x, \Omega \cap \text{Ker}L, 0\} \neq 0.$$

So, condition (b) of Lemma 2.1 is also satisfied. Therefore, from Lemma 2.1 we conclude that system (1) has at least one ω -periodic solution. The proof is complete.

Remark 3.1: If $a_i(t) \equiv b_i(x_i) = b_i x_i$ and $f = g$, then the system (1) reduces to system (2) in [10]. In Theorem 3.1, we remove the continuously differentiable assumptions of the time-varying delays $\tau_{ij}(t)$ and $\sigma_{ij}(t)$.

4. AN ILLUSTRATIVE EXAMPLE

In this section, we give an example to illustrate the effectiveness of our result.

Example 4.1: Consider the following Cohen-Grossberg type neural networks with neutral time-varying delays

$$\begin{cases} \dot{x}_1(t) = -a_1(x_1(t)) [5x_1(t) - 0.5 f_2(x_2(t - \tau_{12}(t))) - 0.4 \sin t \cdot g_1(\dot{x}_1(t - \sigma_{11}(t))) - J_1(t), \\ \dot{x}_2(t) = -a_2(x_2(t)) [4x_2(t) - 0.3 \sin t \cdot f_1(x_1(t - \tau_{21}(t))) - 0.2 g_1(\dot{x}_1(t - \sigma_{21}(t))) - J_2(t), \end{cases} \quad (25)$$

where

$$a_1(u) = 1 + \frac{1}{10} \sin u, \quad a_2(u) = 1 + \frac{1}{5} \sin u, \quad f_1(u) = f_2(u) = \frac{1}{2} \sin u, \quad g_1(u) = \frac{2}{3} u,$$

$$a_{11}(t) = 0, \quad a_{12}(t) = 0.5, \quad a_{21}(t) = 0.3 \sin t, \quad a_{22}(t) = 0, \quad b_{11}(t) = 0.4 \sin t,$$

$$b_{12}(t) = b_{22}(t) = 0, \quad b_{21}(t) = 0.2, \quad J_1(t) = 3 \sin t, \quad J_2(t) = 0.75 \sin t,$$

$\tau_{12}(t)$, $\tau_{21}(t)$, $\sigma_{11}(t)$ and $\sigma_{21}(t)$ can be any positive continuous bounded 2π -periodic functions. Obviously, $f_i(u)$ ($i = 1, 2$) and $g_1(u)$ satisfy the Lipschitz condition (H_4) with constants $L_i = \frac{1}{2}$ and $l_1 = \frac{2}{3}$, respectively. By the direct calculation, we have

$$A = \begin{pmatrix} 5 & -0.25 \\ -0.15 & 4 \end{pmatrix}, \quad B = \begin{pmatrix} 0.2667 & 0 \\ 0.1334 & 0 \end{pmatrix}, \quad C = \begin{pmatrix} 0.7067 & 0 \\ -0.16 & 1 \end{pmatrix},$$

$$D = \begin{pmatrix} 5.5 & 0.275 \\ 0.18 & 4.8 \end{pmatrix}, \quad C^{-1} = \begin{pmatrix} 1.415 & 0 \\ 0.2264 & 1 \end{pmatrix}, \quad A - B(C^{-1}D) = \begin{pmatrix} 2.9244 & -0.3538 \\ -1.1882 & 3.9481 \end{pmatrix},$$

and

$$(A - B(C^{-1}D))^{-1} = \begin{pmatrix} 0.3549 & 0.0318 \\ 0.1068 & 0.2629 \end{pmatrix},$$

which implies that C and $A - B(C^{-1}D)$ are two nonsingular M -matrix. Hence, all the conditions of Theorem 3.1 are satisfied. So, by Theorem 3.1, system (25) has at least one 2π -periodic solution.




Figure 1: Numerical Solution $x_1(t), x_2(t)$ of system (25), where $\tau_{12}(t) = 1$, $\tau_{21}(t) = 0.8$, $\sigma_{11}(t) = 0.6$ and $\sigma_{21}(t) = 0.7$, $x_1(s) = x_2(s) = 0$ for $s \in [-1, 0]$

ACKNOWLEDGEMENTS

Project supported by the Natural Science Foundation of Jiangsu Education Office (06KJB110010) and Jiangsu Planned Projects for Postdoctoral Research Funds.




Figure 2: Phase Trajectories of System (25), where $\tau_{12}(t) = 1$, $\tau_{21}(t) = 0.8$, $\sigma_{11}(t) = 0.6$ and $\sigma_{21}(t) = 0.7$, $x_1(s) = x_2(s) = 0$ for $s \in [-1, 0]$

REFERENCES

- [1] J. Cao, New results concerning exponential stability and periodic solutions of delayed cellular neural networks, *Phys. Lett. A*, Vol. **307**, pp. 136–147, 2003.
- [2] K. Gopalsamy and S. Mohanad, Time delays and stimulus-dependent pattern formation in periodic environments in isolated neurons, *IEEE Trans. Neural Networks*, Vol. **13**, pp. 551–563, 2002.
- [3] S. Guo, L. Huang, Periodic oscillation for a class of neural networks with variable coefficients, *Nonlinear Anal.*, Vol. **6**, pp. 545–561, 2005.
- [4] Y. Li, Existence and stability of periodic solutions for Cohen-Grossberg neural networks with multiple delays, *Chaos, Solitons and Fractals*, Vol. **20**, pp. 459–466, 2004.
- [5] B. Liu and L. Huang, Existence and exponential stability of periodic solutions for cellular neural networks with time-varying delays, *Phys. Lett. A*, Vol. **349**, pp. 474–483, 2006.
- [6] Z. Liu and L. Liao, Existence and global exponential stability of periodic solution of cellular neural networks with time-varying delays, *J. Math. Anal. Appl.*, Vol. **290**, pp. 247–262, 2004.
- [7] Z. Huang and Y. H. Xia, Exponential p -stability of second order Cohen-Grossberg neural networks with transmission delays and learning behavior, *Simulation Modelling Practice and Theory*, Vol. **15**, pp. 622–634, 2007.
- [8] Y. H. Xia, J. Cao and S. S. Cheng, Periodicity in a Lotka Volterra mutualism system with several delays, *Appl. Math. Modelling*, Vol. **31**, pp. 1960–1969, 2007.
- [9] Y. H. Xia and J. Cao, Almost periodic solutions for an ecological model with infinite delays, *Proceeding of Edinburgh Mathematical Society*, Vol. **50**, pp. 229–249, 2007.
- [10] Z. Gui, W. Ge, X. Yang, Periodic oscillation for a Hopfield neural networks with neutral delays, *Phys. Lett. A* Vol. **364**, pp. 267–273, 2007.
- [11] J. Chen, X. Chen, *Special Matrices*, Tsinghua Univ. Press, Beijing, 2001.
- [12] R.E. Gaines, J. L. Mawhin, *Coincidence Degree and Nonlinear Differential Equations*, Springer-Verlag, Berlin, 1977.
- [13] Berman A., Plemmons R. J., *Nonnegative matrices in the mathematical science*, Academic Press, New York, 1979.

-
- [14] Michael Gr. Voskoglou, A Fuzzy Model for Human Reasoning, *International Journal of Mathematics and Engineering with Computer*.
- [15] S.V. Ingale and D. Datta, Evidence Theory Based Uncertainty Quantification in Radiological Risk Due to Accidental Release of Radioactivity from a Nuclear Power Plant, *International Journal Computer Science and System Analysis*.
- [16] Alketa Hyso and Betim Cico, Neural Networks as Improving Tools for Agent Behavior, *International Journal Computer Science and Knowledge Engineering*
- [17] Sasank M., A Goal Programming Model for Measuring the Effectiveness of Quality Control Circles, *International Journal of Mathematics and Applied Statistics*
- [18] Nicholas M, Specification and Informational Issues in Credit Scoring, *Journal of Statistical Sciences*.
- [19] Basim Makhool., Access to Finance and the Potential Impact of New Collateral Schemes on Palestinian SMEs Financial Needs, *The Global Journal of Finance and Economics*, Vol. 4, No. 1, pp. 1-12.
- [20] Qiqing Yu, Buckley–James–type estimators in classical–cohort studies, *International Journal of Business Statistics and Finance*.
- [21] G. Narsimlu and L. Anand Babu, MHD Free Convection Flow in Porous Medium between Two Heated Flat Vertical Plates, *International Journal of Scientific Computing*.
- [22] Maged George Iskander, A Bi-Criteria Approach for Chance-Constrained Goal Programming Under Uncertainty: A Computational Study, *International Review of Fuzzy Mathematics*.
- [23] K. Karthikeya, Existence and Uniqueness of a Fuzzy Impulsive Solution for the Nonlinear Fuzzy Impulsive Neutral Functional Differential Equations, *International Journal of Fuzzy Systems and Rough Systems*.
- [24] Jian-Wei LIU, Research on Traffic Data-Collecting System Based on MC9S12D64 Microcontroller, *Journal of Information Technology and Engineering*
- [25] Halime Temel Nalin, Impact of Oil Price Shocks on Macroeconomic Variables: Evidence from an Emerging Market, *Global Review of Business and Economic Research*
- [26] Govindappa Navalagi, Ratnesh Saraf, and Miguel Caldas, On Weak Forms of Fuzzy Semi-preopen and Fuzzy Semi-preclosed Functions, *International Journal of Engineering Computer Science and Mathematics*.
- [27] A. S. K. Ratnam and M. Venkata Rao, Adaptive Fault Tolerant Routing in Interconnection Networks: A Review, *International Journal of Mathematics Computer Sciences and Information Technology*.
- [28] S. Lakshmi and R. Balakumar, A Stochastic Model for Finding the Survival Function of PRA, PA and Plasma ANP of the Human Subjects by using BTLG Model, *Global Journal of Applied Mathematics and Mathematics Sciences*.
- [29] Patricia Mariela Morillas, Orthogonal Wavelet Vectors and Multiresolution Analysis In Hilbert Spaces, *International Journal of Mathematical Sciences*.
- [30] M. M. El-Dessoky, A Simple Global Synchronization for Coupled Chaotic System, *International Journal of Applied Mathematics and Applications*.
- [31] Raed I. Hamed, Towards a Formal Fuzzy Framework for Modeling and Stationary Analysis of GRNs, *International Journal of Computer Science and Management System*.
- [32] Ha Nguyen, A New Method for Calculating Weight of Sentence Based on Amount of Information and Linguistic Score, *International Journal of Advanced Computer Engineering*.
- [33] Dayal R.P, A Study of Various Methodologies used for Navigation of Autonomous Mobile Robot, *International Journal of Applied Artificial Intelligence in Engineering System*.
- [34] Rajinder Tiwari and R. K. Singh, An Overview of the Technical Development of the Current Mirror used in Analog CMOS Circuits, *International Journal of Computer and Electronics Engineering*
- [35] Panayotis Fouliras, Using DEA for Network Efficiency Evaluation, *International Journal of Engineering and Management, International Journal of Engineering and Management*
- [36] Umesh Sehgal, Performance Analysis & Enhancements in Intrusion Detection System for Wireless Local Area Networks, *International Journal of Applied Computing*

- [37] Asha Sharma., Cost of Capital and Its Impact on Profitability in Indian Industries, *International Journal of Business Management Economics and Information Technology*, Vol. 4, No. 2, Pp. 157-165.
- [38] Sonja B. Langley & Alison M. Keefe., An Analysis of UK Net Beef Imports during the Mad Cow Crisis: Implications of a J-Shaped Curve, *International Economics and Finance Journal*, Vol. 1, No. 2, p. 275.
- [39] Takashi Obara., Fluctuation-Dissipation Theorems on Foreign Exchange Data, *International Journal of Financial Economics and Econometrics*, Vol. 1, No. 1, pp 1-9.
- [40] Ladislaus M , Indigenous Knowledge, Postcolonialism and Indigenous Education Politics in African Perspective, *JOURNAL OF SOCIAL ANTHROPOLOGY*
- [41] Laura C. Rivero, Equality Dependencies: Analysis, Modeling and Conceptual Issues, *International Journal of Computer Science and Software Technology*
- [42] Jun-Xiang Wang and Zhe-Ming Lu, A High Capacity Lossless Data Hiding Scheme Based on VQ Index Residual Coding, *International Journal of Information Analysis and Processing*
- [43] Sourav Pal, Research Opportunities for Video Streaming in Kalyan Basu, Sumantra R. Kundu & Wireless Network, *International Computer Science and Information Engineering*
- [44] Kai-Lin Tian, A Reliable Anomaly Detector against Low-rate DDoS Attack, *International Journal of Electronics and Computers*
- [45] Kiran Reddy, Newtcp A Refined New Model For Efficient Explicit Congestion Reduction In High Traffic High Speed Networks Through Automated Rate Controlling, *International Journal of Information Technology and Knowledge Management*.
- [46] Ajay Goe, A New Cryptography Based Watermarking Technique for Information Security, *International Journal of Electronics Engineering*.
- [47] Nicola Costantino & Roberta Pellegrino., Real Option Approach for the Evaluation of Investment Projects The Case of a Multi-Fuel Power Station, *International Journal of Engineering and Management*, Vol. 1, No. 1, pp. 47-53.
- [48] K. Nikolopoulos, D. Thomakos & F. Petropoulos, A. Litsa, V. Assimakopoulos., Forecasting S&P 500 with the Theta Model, *International Journal of Financial Economics and Econometrics*, Vol. 4, No. 1, pp. 73-78
- [49] C. D. Lai. Hazard Rate that Asymptotes to a Constant and Applications, *International Journal of Reliability and Quality Performance*, Vol. 6 no. 1 pp 1-9.
- [50] Alan Whaley , William E. Gillis and Ermanno Affuso., A Panel Data Analysis of Efficiency and Productivity of the U.S. Hospital Care System; *Review of Applied Economics*, Vol. 13 nos. 1-2 pp 1-20.
- [51] Spyridon D. Symeonides.; On the Finite-Sample Distribution of the Feasible GLS Estimator: An Application of the Cornish-Fisher Correction; *The Journal of World Economic Review*, Vol. 12 no. 1; pp 1-13.
- [52] GEORGE SOKLIS.; Labour versus Alternative Value Bases in Actual Joint Production Systems, *Bulletin of Political Economy*, Vol. 9:1 pp1-31.
- [53] Ali A.L. Kilindo.; Does Public Investment Determine Private Investment? A Multivariate Cointegration Analysis of Public/Private Investment Linkages In Tanzania; *Asian-African Journal of Economics and Econometrics*, Vol. 16 no. 1 pp 1-11.
- [54] Ran Zhang (2017); Does Systematic Risk Affect A-H Price Spread?; *Journal of Money Banking and Finance*, Vol. 1 no. 1 pp 1-11.
- [55] Hab. W.Szymanowski & Nadzw. SGGW (2017); Competitive Strategies for Future Supply Chain Management in European Union on the Food Example; *Global Journal of Accounting and Economic Research*, Vol. 1 no. 1; pp 1-16.