

**SOLUTIONS TO ABC-FRACTIONAL ORDER NEUTRAL
MIXED INTEGRO DELAY DIFFERENTIAL EQUATIONS WITH
IMPULSE**

R. DEVIPRIYA, S. SELVI, KOTTAKKARAN SOOPPY NISAR*, AND C. RAVICHANDRAN

ABSTRACT. This work establishes the existence and uniqueness (EUS) of solutions for impulsive neutral mixed integro fractional delay differential equations involving recently explored ABC-fractional derivatives. Fixed-point techniques are applied to confirm EUS of solutions of ABC- fractional order integrodifferential system with impulse.

1. Introduction

Fractional calculus is a generalised classical calculus with order of real or complex values. In recent decades, it has developed tremendously and has turned into one of the most powerful tools to describe dynamical systems in numerous scientific disciplines. Unlike the ordinary differential operator, fractional derivatives are global in nature and produce concise results by using minimum parameters than that of the classical model. For the fundamental developments in this field, one can refer [41, 44]. Because of these astonishing results and more flexibility, some of the researchers not only exploited fractional operators in modelling and understanding the complex system, but also enriched the calculus with various fractional operators. Some of the fractional operators contain singular kernel. To prevail over this, Caputo and Fabrizio defined an operator involving an exponential function which is non-singular[17]. Many researchers have worked on developing the theory of FDE using CF derivative [1, 3, 11, 26, 35, 37, 47, 48].

Motivated by CF derivative, Atangana and Baleanu (AB) put forward a novel nonsingular fractional derivative possessing the Mittag-Leffler function and it influences new attributes [12]. Many researchers have tried to replace the ordinary differential operator with AB fractional derivative in physical systems, as it is more applicable and yields effective results [4, 5, 6, 7, 13, 23, 24, 28, 29, 30, 31, 33, 34, 38, 39, 40, 42, 43, 50, 51]. Alkahtani et al.[10] did numerical research on dynamics of Chau's loop. Gomez et al. [22] obtained analytical solutions for the electric loop with AB derivatives. Chang et al. [54] applied AB derivatives to solve variational Herglotz problems. Kamal Shah et al. [46] analyzed the qualitative behaviour of the mathematical model of SARS - COV -19 under fractional non-singular derivative.

2000 *Mathematics Subject Classification.* 34A08; 34K40; 58C30.

Key words and phrases. ABC-derivatives; Fractional differential equations; Fixed point theorem.

*Corresponding author.

Some dynamical problems subjected to short-term perturbations, which allow discontinuities in the evolution phenomena, are studied using impulsive fractional differential equations.[15, 16, 18, 19, 25, 36]. we refer the monographs [14, 32, 45]. Mathematical models involving the fractional differential operators with delay, depend on duration of certain hidden processes as past history [2, 8, 21].Integro-differential equations are generalizations of FDE, Fredholm and Volterra integral equations and are approximation to partial differential equations, which represent much of the continuum phenomena. These kinds of equations are employed in modeling many problems in mathematical physics, biophysics, rheology such as heat conduction in materials with memory.[9, 20, 27, 52, 53, 55].

Hasib Khan et al.[28] ensured EU and data dependence of the solution of impulsive FDE with ABC-derivative using Schauder fixed point theorem. Recently, Ravichandran et al.[43] proved the EUS results of neutral impulsive FIDE's in the ABC sense of the form:

$$\begin{aligned} {}_0^{ABC}D^\gamma[v(t) - \Theta(t, v_t)] &= f\left(t, v_t, \int_0^t k(t, s, v_s)ds\right), \quad 0 < \gamma < 1 \\ \Delta v(t_k) &= v(t_k^+) - v(t_k^-) = I_k^* v(t_k), \\ v|_{[-\tau^*, 0]} &= v_0, \end{aligned}$$

where $v(t) \in C[0, 1]$, $f : \mathcal{J} \times PC^1 \times \mathcal{J} \rightarrow \mathbb{R}^n$ is continuous. $I_k^* : \mathbb{R}^n \rightarrow \mathbb{R}^n$, $v(t_s) = v(t+s)$ for $-\tau^* \leq s \leq 0$ where $v(t_k^+) = \lim_{\delta \rightarrow 0^+} v(t_k + \delta)$ and $v(t_k^-) = \lim_{\delta \rightarrow 0^-} v(t_k - \delta)$.

In papers [25], [38] and other, the existence of non linear FDE is proved using Schuader and Krasnoselskii fixed point theorem, whereas in this manuscript, the existence of Neutral FIDE of mixed type with impulses and delay is discussed using Schaefer's fixed point theorem and the obtained results are verified with an example. Inspired by the above works [28] and [43], we have presented the EU of the solution for the following impulsive ABC- fractional neutral mixed type integrodifferential equation using Schaefer's fixed point theorem,

$${}_0^{ABC}D^v \left[x(t) - \sigma(t, x_t) \right] = \mathcal{B}\left(t, x_t, \int_0^t \eta(t, s, x_s)ds, \int_0^T \xi(t, s, x_s)ds\right) \quad (1.1)$$

$$t \in \mathcal{J}^c = \mathcal{J} - \{t_1, t_2, \dots, t_m\}, \quad \mathcal{J} = [0, T], \quad x \neq t_k$$

$$\Delta x(t_k) = x(t_k^+) - x(t_k^-) = I_k^* x(t_k), \quad (1.2)$$

$$x|_{[-\tau^*, 0]} = \zeta + \mathcal{F}^*.$$

where $v \in (0, 1)$, $\mathcal{F}^*(x) = \sum_{i=1}^m \lambda_i x_i(t)$, $x_i \in PC^1$, $\sum_{i=1}^m \lambda_i < 1$ for $i = 1, 2, \dots, m$, $\mathcal{B}\left(t, x_t, \int_0^t \eta(t, s, x_s)ds, \int_0^T \xi(t, s, x_s)ds\right)$ and $\sigma(t, x_t)$ are Lebesgue measurable functions such that $\mathcal{B} \in C\left(\mathcal{J} \times X \times X \times X \rightarrow X\right)$ where X is a Banach space X, $\sigma : \mathcal{J} \times PC^1 \rightarrow \mathbb{R}^n$ are piecewise continuous and $I_k^* : \mathbb{R}^n \rightarrow \mathbb{R}^n$ are continuous on PC^1 where $PC^1 = PC^1\left([-\tau^*, 0], \mathbb{R}^n\right)$ is the space of piecewise continuous function with norm $\|x\|_\infty = \sup_{t \in [-\tau^*, T]} \|x(t)\|$ and $\zeta : [-\tau^*, 0] \rightarrow \mathbb{R}^n$. $\eta, \xi : \mathcal{D} \times X \rightarrow X$

are continuous, where $\mathcal{D} = \{(x, s) \in \mathcal{J} \times \mathcal{J} : t \geq s\}$. Then $x_t(s) = x(t+s)$ for $-\tau^* < s \leq 0$, $x(t_k^+) = \lim_{\delta \rightarrow 0^+} x(t_k + \delta)$ and $x(t_k^-) = \lim_{\delta \rightarrow 0^-} x(t_k - \delta)$.

Consider $Px(t) = \int_0^t \eta(t, s, x_s)ds$ and $Qx(t) = \int_0^T \xi(t, s, x_s)ds$. Then (1.1) becomes

$$\begin{aligned} {}_0^{ABC}D^v \left[x(t) - \sigma(t, x_t) \right] &= \mathcal{B} \left(t, x_t, Px(t), Qx(t) \right), \\ t \in \mathcal{J}^c &= \mathcal{J} - \{t_1, t_2, \dots, t_m\}, \quad \mathcal{J} = [0, T], \quad x \neq t_k, \\ \Delta x(t_k) &= x(t_k^+) - x(t_k^-) = I_k^* x(t_k), \\ x|_{[-\tau^*, 0]} &= \zeta + \mathcal{F}^*. \end{aligned} \tag{1.3}$$

The aim of this paper is to obtain EUS of the above IVP of ABC-fractional order impulsive neutral mixed integrodifferential equation with delay using Schaefer's fixed point theorem.

The rest of the paper is sorted out as follows: In Section 2, definitions and lemmas of ABC-fractional derivatives are recalled. In Section 3, integral form of ABC-derivative of the above-mentioned problem is proved. In Section 4, the main result - EUS of the proposed ABC-fractional order impulsive neutral differential equation is constructed. In the last section, the obtained results are verified with an example.

2. Prerequisite

In this segment, we elicit definitions and tools on fractional operators with the non-singular kernel[44, 41, 17, 12, 35].

For $0 < \alpha < 1$, the Caputo fractional derivative of order α starting at a is given as

$${}_0^C D^\alpha y(t) = \frac{1}{\Gamma(1-\alpha)} \int_a^t (t-s)^{-\alpha} y'(s) ds.$$

Let $\psi \in H^*(a, b)$, $a < b$, v in $[0, 1]$. The ABC - fractional derivative of ψ of order v is defined by

$${}_0^{ABC} D^v \psi(\tau^*) = \frac{B^*(v)}{1-v} \int_0^{\tau^*} \psi'(s) E_v \left[\frac{-v(\tau^* - s)^v}{1-v} \right] ds$$

where $B^*(0) = B^*(1) = 1$.

The associated AB fractional -integral of a function ψ is defined by

$${}_0^{AB} I_t^v \psi(\tau^*) = \frac{1-v}{B^*(v)} \psi(\tau^*) + \frac{v}{B^*(v)\Gamma(v)} \int_0^{\tau^*} \psi(s) (\tau^* - s)^{v-1} ds.$$

Lemma 2.1. *Newton -Leibntiz formula holds for the ABC-fractional order differential and integral operators.*

$${}_0^{AB} I_t^v \left({}_0^{ABC} D_{\tau^*}^v (\theta) \right) = \psi(\tau^*) - \psi(0)$$

For $\psi(t)$ defined on $[e, f]$ and $v \in (m, m + 1)$ for some $m \in N_0$,

$$\begin{aligned} {}_e^{ABR} D^v \left({}_e^{AB} I^v \psi(t) \right) &= \psi(t) \\ {}_e^{AB} I^v \left({}_e^{ABR} D^v \psi(t) \right) &= \psi(t) - \sum_{k=0}^{m-1} \psi^{(k)}(e) \frac{(t-e)^k}{\Gamma(k-1)}. \end{aligned}$$

Theorem 2.2. (Schaefer's fixed point theorem) [49]: A continuous and compact operator ϕ has a fixed point, if the set $\{u \in X; u = \lambda\phi u \text{ for } 0 \leq \lambda \leq 1\}$ is bounded.

3. Integral form of ABC - FIDE

Here, we analyse the impulsive neutral ABC - FIDE and its integral form.

Theorem 3.1. For $v \in (0, 1)$ and $\mathcal{B}, \sigma \in PC^1[0, T]$ such that $\sigma(0, x_0) = 0$ and $\mathcal{B}\left(0, x_0, 0, \int_0^T \xi(0, s, x_s) ds\right) = 0$, $x(t)$ is a solution of (1.3) provided that

$$x(t) = \begin{cases} \zeta + \mathcal{F}^*, & t \in [-\tau^*, 0] \\ \zeta(0) + \mathcal{F}^*(0) + \sigma(t, x_t) + {}_0^{AB} I^v \mathcal{B}\left(t, x_t, Px(t), Qx(t)\right), & t \in [0, t_1] \\ \zeta(0) + \mathcal{F}^*(0) + \sigma(t, x_t) + {}_0^{AB} I^v \mathcal{B}\left(t, x_t, Px(t), Qx(t)\right) \\ + I_1^*(x(t_1^-)), & t \in [t_1, t_2] \\ \cdot \\ \cdot \\ \zeta(0) + \mathcal{F}^*(0) + \sigma(t, x_t) + {}_0^{AB} I^v \mathcal{B}\left(t, x_t, Px(t), Qx(t)\right) \\ + \sum_{k=1}^m I_k^*(x(t_k^-)). & t \in [t_m, T] \end{cases}$$

Proof. The solution $x(t)$ of the problem (1.3), we have for $t \in [0, t_1]$,

$$\begin{aligned} {}_0^{AB} I^v \left[{}_0^{ABC} D^v \left\{ x(t) - \sigma(t, x_t) \right\} \right] &= {}_0^{AB} I^v \left[\mathcal{B}\left(t, x_t, Px(t), Qx(t)\right) \right] \\ x(t) - \sigma(t, x_t) &= {}_0^{AB} I^v \left[\mathcal{B}\left(t, x_t, Px(t), Qx(t)\right) \right] + c_0 \\ x(t) &= \zeta(0) + \mathcal{F}^*(0) + \sigma(t, x_t) + {}_0^{AB} I^v \left[\mathcal{B}\left(t, x_t, Px(t), Qx(t)\right) \right] \end{aligned}$$

and

$$x(t_1^-) = \zeta(0) + \mathcal{F}^*(0) + \sigma(t_1, x_{t_1}) + {}_0^{AB} I^v \left[\mathcal{B}\left(t_1, x_{t_1}, Px(t_1), Qx(t_1)\right) \right].$$

For $t \in (t_1, t_2]$, $\Delta x(t_k) = I_k^* x(t_k)$, we have $x(t_1^+) - x(t_1^-) = I_1^* x(t_1)$

$$\begin{aligned} x(t) &= x(t_1^+) - x(t_1) + x(t) \\ &= x(t_1^-) + I_1^*(x(t_1^-)) - \sigma(t_1, x_{t_1}) - \zeta(0) - \mathcal{F}^*(0) - \\ &\quad {}_0^{AB} I^v \left[\mathcal{B}\left(t_1, x_{t_1}, Px(t_1), Qx(t_1)\right) \right] + \zeta(0) + \mathcal{F}^*(0) + \sigma(t, x_t) \\ &\quad + {}_0^{AB} I^v \left[\mathcal{B}\left(t, x_t, Px(t), Qx(t)\right) \right] \end{aligned}$$

Using $x(t_1^-)$ in the above equation, we get

$$x(t) = \zeta(0) + \mathcal{F}^*(0) + \sigma(t, x_t) + I_1^*(x(t_1^-)) + {}_0^{AB} I^v \left[\mathcal{B}\left(t, x_t, Px(t), Qx(t)\right) \right].$$

For $t \in (t_2, t_3]$, we have

$$\begin{aligned} x(t) &= x(t_2^+) - x(t_2) + x(t) \\ &= x(t_2^-) + I_2^*(x(t_1^-)) - \sigma(t_2, x_{t_2}) - \zeta(0) - \mathcal{F}^*(0) \\ &\quad - {}_0^{AB} I^v \left[\mathcal{B}(t_2, x_{t_2}, Px(t_2), Qx(t_2)) \right] + \zeta(0) + \mathcal{F}^*(0) + \sigma(t, x_t) \\ &\quad + {}_0^{AB} I^v \left[\mathcal{B}(t, x_t, Px(t), Qx(t)) \right]. \end{aligned}$$

Using $x(t_2^-)$ in the above equation, we get

$$x(t) = \zeta(0) + \mathcal{F}^*(0) + \sigma(t, x_t) + \sum_{k=1}^2 I_k^*(x(t_k^-)) + {}_0^{AB} I^v \left[\mathcal{B}(t, x_t, Px(t), Qx(t)) \right].$$

As we proceed, it leads to the case when $t \in (t_m, T]$ and we get

$$x(t) = \zeta(0) + \mathcal{F}^*(0) + \sigma(t, x_t) + \sum_{k=1}^m I_k^*(x(t_k^-)) + {}_0^{AB} I^v \left[\mathcal{B}(t, x_t, Px(t), Qx(t)) \right].$$

□

4. Theorems for Main Results

Let $h : \mathcal{J} \rightarrow \mathbb{R}$ is measurable in $\mathcal{L}^\omega(\mathcal{J})$ with the norm:

$$\|h\|_{\mathcal{L}}^\omega = \begin{cases} \left(\int_{\mathcal{J}} |g(t, x_t)|^\omega dt \right)^{\frac{1}{\omega}}, & 1 \leq \omega < \infty \\ \inf_{\mu(\bar{J})} \left(\sup_{t \in J - \bar{J}} |g(t, x_t)| \right), & \omega = \infty \end{cases}$$

where $\mu(\bar{J})$ is the Lebesgue measure on \bar{J} and $\|h\|_{\mathcal{L}}^\omega < \infty$. Now we define an operator Ω as

$$\Omega x(t) = \begin{cases} \zeta + \mathcal{F}^*, & t \in [-\tau^*, 0] \\ \zeta(0) + \mathcal{F}^*(0) + \sigma(t, x_t) + {}_0^{AB} I^v \mathcal{B}(t, x_t, Px(t), Qx(t)), & t \in [0, t_1] \\ \zeta(0) + \mathcal{F}^*(0) + \sigma(t, x_t) + {}_0^{AB} I^v \mathcal{B}(t, x_t, Px(t), Qx(t)) \\ + I_1^*(x(t_1^-)), & t \in [t_1, t_2] \\ \cdot \\ \cdot \\ \zeta(0) + \mathcal{F}^*(0) + \sigma(t, x_t) + {}_0^{AB} I^v \mathcal{B}(t, x_t, Px(t), Qx(t)) \\ + \sum_{k=1}^m I_k^*(x(t_k^-)). & t \in [t_m, T] \end{cases}$$

For the proof of our main result, we need the following assumptions

- (A1) Let $x \in \mathcal{PC}^1[0, T]$ and suppose that $\mathcal{B} \in C(\mathcal{J} \times X \times X \times X, X)$ is piecewise continuous and there exists positive constants $\mathcal{L}_1, \mathcal{L}_2$ and \mathcal{L} such that
- $$\|\mathcal{B}(t, x_1, u_1, z_1) - \mathcal{B}(t, x_2, u_2, z_2)\| \leq \mathcal{L}_1 \left(\|x_1 - x_2\| + \|u_1 - u_2\| + \|z_1 - z_2\| \right)$$
- for all x, u, z in Y , where $Y = C[J, X]$ be the set of continuous functions

on J with values in the Banach space X , $\mathcal{L}_2 = \max_{t \in \mathcal{J}} \|\mathcal{B}(t, 0, 0, 0)\|$ and $\mathcal{L} = \max\{\mathcal{L}_1, \mathcal{L}_2\}$.

- (A2) There exists positive constants $\mathbb{N}_1, \mathbb{N}_2$ and \mathbb{N} such that
 $\|P(t, s, x_1) - P(t, s, x_2)\| \leq \mathbb{N}_1(\|x_1 - x_2\|)$
for all x_1, x_2 in Y , $\mathbb{N}_2 = \max_{(t,s) \in \mathcal{D}} \|P(t, s, 0)\|$ and $\mathbb{N} = \max\{\mathbb{N}_1, \mathbb{N}_2\}$.
- (A3) There exists positive constants $\mathbb{C}_1, \mathbb{C}_2$ and \mathbb{C} such that
 $\|Q(t, s, x_1) - Q(t, s, x_2)\| \leq \mathbb{C}_1(\|x_1 - x_2\|)$
for all x_1, x_2 in Y , $\mathbb{C}_2 = \max_{(t,s) \in \mathcal{D}} \|Q(t, s, 0)\|$ and $\mathbb{C} = \max\{\mathbb{C}_1, \mathbb{C}_2\}$.
- (A4) Let the impulses $I_k^* \in \mathcal{C}(\mathcal{R}^n, \mathcal{R}^n)$ be bounded and for a $\vartheta^* > 0$, we have
 $\mu(t) \in L^{\frac{1}{\delta}}(J)$ with $\left\| I_k^*(x(t_k^-)) - I_k^*(u(t_k^-)) \right\| \leq \vartheta^*(t) \|x - u\|_{\infty}$
for each $x, u \in PC^1([- \tau^*, T])$ (ζ), and for each $r \in I^+$, $k = 1, 2, \dots, m$
then $\mathcal{N}^* = \max \left\{ \left\| I_k^*(x(t_k^-)) \right\| : \|x\| \leq r \right\}$.
- (A5) Let for a $\delta \in (0, h)$ and there exists function $\mu(t)$ in $\mathcal{L}^{\frac{1}{\delta}}(\mathcal{J})$ such that
 $\|\sigma(t, x_t) - \sigma(t, u_t)\| \leq \mu(t) \|x_t - u_t\|$, for $t \in \mathcal{J}$ and $x_t, u_t \in \mathcal{PC}^1$.

Lemma 4.1. *If (A1) – (A3) are satisfied, then for any $t \in \mathcal{J}$ and $x_1, x_2 \in Y$,
 $\|Px_1(t) - Px_2(t)\| \leq \mathbb{N}t \|x_1 - x_2\|$, $\|Qx_1(t) - Qx_2(t)\| \leq \mathbb{C}T \|x_1 - x_2\|$,
 $\|Px(t)\| \leq t(\mathbb{N}_1 \|x\| + \mathbb{N}_2)$, $\|Qx(t)\| \leq T(\mathbb{C}_1 \|x\| + \mathbb{C}_2)$.*

Theorem 4.2. *: If the conditions (A1) – (A5) hold, and $(\mathbb{L} \|x_{t_2} - x_{t_1}\| + \mathbb{N}t \|x_{t_2} - x_{t_1}\| + \mathbb{C}T \|x_{t_2} - x_{t_1}\|) = q(t_2 - t_1)$ and $q \in \mathcal{L}^1(\mathcal{J}, \mathcal{R}_+)$ and if the inequality*

$$\left[\|\zeta(0) + \mathcal{F}^*(0)\| + \|\sigma(t, x_t)\| + m\mathcal{N}^* + \left(\frac{1-v}{B^*(v)} + \frac{t^v}{B^*(v)\Gamma v} \right) p \|x_t\| \right] \leq \varrho^* \quad (4.1)$$

holds, then the problem (1.3) has a solution where $p \|x_t\| = (\mathcal{L} \|x_t\| + \mathbb{N}t \|x_t\| + \mathbb{C}T \|x_t\|)$ and for each positive r , $\mathbb{B}_r \in \{x \in Y : \|x\| \leq r\}$, $k = 1, 2, \dots, m$.

Proof. We divide the proof in the following parts.

Part 1: Ω is continuous

Let $\{x_n\}$ be a sequence such that $x_n \rightarrow x$ in $PC^1([- \tau^*, T])$.

Since $\mathcal{B}(t, x_t, Px(t), Qx(t))$, $\sigma(t, x_t)$ and the impulses $I_k^*(x(t_k^-))$ are continuous operators on $PC^1([- \tau^*, \mathbb{R}^n])$ (ζ) for $k = 1, 2, \dots, m$. By using (A1) and lemma

4.2, then for each $t \in [0, t_1]$, we have

$$\begin{aligned}
 \|\Omega x_n - \Omega x\| &\leq \|\sigma(t, x_{n_t}) - \sigma(t, x_t)\| + \left\| {}_0^{AB} I^v \mathcal{B}(t, x_{n_t}, Px_n(t), Qx_n(t)) \right. \\
 &\quad \left. - \mathcal{B}(t, x_t, Px(t), Qx(t)) \right\| \\
 &\leq \mu(t)\|x_n - x\| + \frac{1-v}{B^*(v)} \left\| \mathcal{B}(t, x_{n_t}, Px_n(t), Qx_n(t)) \right. \\
 &\quad \left. - \mathcal{B}(t, x_t, Px(t), Qx(t)) \right\| + \frac{v}{B^*(v)} \left\| I_0^v \mathcal{B}(t, x_{n_t}, Px_n(t), Qx_n(t)) \right. \\
 &\quad \left. - I_0^v \mathcal{B}(t, x_t, Px(t), Qx(t)) \right\| \\
 &\leq \mu(t)\|x_n - x\| + \frac{1-v}{B^*(v)} (\mathbb{L}\|x_n - x\| + \mathbb{N}t\|x_n - x\| + \mathbb{C}T\|x_n - x\|) \\
 &\quad + \frac{v}{B^*(v)} (\mathbb{L}\|x_n - x\| + \mathbb{N}t\|x_n - x\| + \mathbb{C}T\|x_n - x\|) I_0^v(t) \\
 &\leq \mu(t)\|x_n - x\| + \left[\frac{1-v}{B^*(v)} + \frac{t^v}{B^*(v)\Gamma v} \right] p\|x_n - x\|
 \end{aligned}$$

$\|\Omega x_n - \Omega x\| \rightarrow 0$ as $x_n \rightarrow x$.

Similarly for $t \in (t_1, t_2]$, by using (A4) and (A5), we have

$$\begin{aligned}
 \|\Omega x_n - \Omega x\| &\leq \|\sigma(t, x_{n_t}) - \sigma(t, x_t)\| + \left\| {}_0^{AB} I^v \mathcal{B}(t, x_{n_t}, Px_n(t), Qx_n(t)) \right. \\
 &\quad \left. - \mathcal{B}(t, x_t, Px(t), Qx(t)) \right\| + \|I_1 x_n(t_1^-) - I_1 x(t_1^-)\| \\
 &\leq \mu(t)\|x_n - x\| + \frac{1-v}{B^*(v)} \left\| \mathcal{B}(t, x_{n_t}, Px_n(t), Qx_n(t)) \right. \\
 &\quad \left. - \mathcal{B}(t, x_t, Px(t), Qx(t)) \right\| + \frac{v}{B^*(v)} \left\| I_0^v \mathcal{B}(t, x_{n_t}, Px_n(t), Qx_n(t)) \right. \\
 &\quad \left. - I_0^v \mathcal{B}(t, x_t, Px(t), Qx(t)) \right\| + \vartheta^* \|x_n - x\| \\
 &\leq \left[\mu(t) + \vartheta^*(t) \right] \|x_n - x\| + \frac{1-v}{B^*(v)} (\mathbb{L}\|x_n - x\| + \mathbb{N}t\|x_n - x\| \\
 &\quad + \mathbb{C}T\|x_n - x\|) + \frac{v}{B^*(v)} (\mathbb{L}\|x_n - x\| + \mathbb{N}t\|x_n - x\| \\
 &\quad + \mathbb{C}T\|x_n - x\|) I_0^v(t) \\
 &\leq \left[\mu(t) + \vartheta^*(t) \right] \|x_n - x\| + \left[\frac{1-v}{B^*(v)} + \frac{t^v}{B^*(v)\Gamma v} \right] p\|x_n - x\|
 \end{aligned}$$

Hence $\|\Omega x_n - \Omega x\| \rightarrow 0$ as $x_n \rightarrow x$. Ultimately for $t \in (t_m, T]$, we have

$$\begin{aligned}
 \|\Omega x_n - \Omega x\| &\leq \|\sigma(t, x_{n_t}) - \sigma(t, x_t)\| + \left\| {}_0^{AB} I^v \mathcal{B}(t, x_{n_t}, Px_n(t), Qx_n(t)) \right. \\
 &\quad \left. - {}_0^{AB} I^v \mathcal{B}(t, x_t, Px(t), Qx(t)) \right\| + \sum_{k=1}^m \left\| I_1^*(x_n(t_1^-)) - I_1^*(x(t_1^-)) \right\|
 \end{aligned}$$

$$\leq \left[\mu(t) + m\vartheta^*(t) \right] \|x_n - x\| + \left[\frac{1-v}{B^*(v)} + \frac{t^v}{B^*(v)\Gamma(v)} \right] p \|x_n - x\|$$

$\|\Omega x_n - \Omega x\| \rightarrow 0$ as $n \rightarrow \infty$. Hence, Ω is continuous.

Part 2: Ω is bounded in $PC^1([- \tau^*, T])$

It is enough to show that for any $\delta^* > 0$, then $x \in \mathbb{B}_{\delta^*} = \{x \in PC^1([- \tau^*, T], \mathbb{R}^n)\}; \|x\| \leq \delta^*\}$. By using the inequality (4.1) and lemma 4.2, for each $t \in [0, t_1]$,

$$\begin{aligned} \|\Omega x(t)\| &\leq \|\zeta(0) + \mathcal{F}^*(0)\| + \|\sigma(t, x_t)\| + \frac{1-v}{B^*(v)} \left\| \mathcal{B}(t, x_t, Px(t), Qx(t)) \right\| \\ &\quad + \frac{v}{B^*(v)} \left\| \mathcal{B}(t, x_t, Px(t), Qx(t)) \right\| I_0^v(t) \\ &\leq \|\zeta(0) + \mathcal{F}^*(0)\| + \|\sigma(t, x_t)\| + \left[\frac{1-v}{B^*(v)} \right. \\ &\quad \left. + \frac{t^v}{B^*(v)\Gamma(v)} \right] (\mathbb{L}\|x_t\| + \mathbb{N}t\|x_t\| + \mathbb{C}T\|x_t\|) \\ &\leq \|\zeta(0) + \mathcal{F}^*(0)\| + \|\sigma(t, x_t)\| + \left[\frac{1-v}{B^*(v)} + \frac{t^v}{B^*(v)\Gamma(v)} \right] p \|x_t\| \\ &\leq \varrho^*. \end{aligned}$$

$\|I^y$, by using (A4), for each $t \in (t_m, T]$.

$$\begin{aligned} \|\Omega t(t)\| &\leq \|\zeta(0) + \mathcal{F}^*(0)\| + \|\sigma(t, x_t)\| + \frac{1-v}{B^*(v)} \left\| \mathcal{B}(t, x_t, Px(t), Qx(t)) \right\| \\ &\quad + \frac{v}{B^*(v)} \left\| \mathcal{B}(t, x_t, Px(t), Qx(t)) \right\| I_0^v(t) + \sum_{k=1}^m \left\| I_k^*(x(t_k^-)) \right\| \\ &\leq \|\zeta(0) + \mathcal{F}^*(0)\| + \|\sigma(t, x_t)\| \\ &\quad + \left[\frac{1-v}{B^*(v)} + \frac{t^v}{B^*(v)\Gamma(v)} \right] (\mathbb{L}\|x_t\| + \mathbb{N}t\|x_t\| + \mathbb{C}T\|x_t\|) + m\mathcal{N}^* \\ &\leq \|\zeta(0) + \mathcal{F}^*(0)\| + \|\sigma(t, x_t)\| + \left[\frac{1-v}{B^*(v)} + \frac{t^v}{B^*(v)\Gamma(v)} \right] p \|x_t\| + m\mathcal{N}^* \\ &\leq \varrho^*. \end{aligned}$$

This proves that Ω is bounded.

Part 3: Ω is equicontinuous in $PC^1([- \tau^*, T], \mathbb{R}^n)$.

Let $t_1, t_2 \in (0, t_1]$

$$\begin{aligned} \|\Omega x(t_2) - \Omega x(t_1)\| &\leq \frac{1-v}{B^*(v)} \left\| \mathcal{B}(t_2, x_{t_2}, Px(t_2), Qx(t_2)) \right. \\ &\quad \left. - \mathcal{B}(t_1, x_{t_1}, Px(t_1), Qx(t_1)) \right\| \end{aligned}$$

$$\begin{aligned}
 & + \frac{v}{B^*(v)} \left[\left\| \mathcal{B}(t_2, x_{t_2}, Px(t_2), Qx(t_2)) I_0^v(t_2) \right. \right. \\
 & \quad \left. \left. - \mathcal{B}(t_1, x_{t_1}, Px(t_1), Qx(t_1)) I_0^v(t_1) \right\| \right] \\
 & + \|\sigma(t_2, x_{t_2}) - \sigma(t_1, x_{t_1})\| \\
 & \leq \frac{1-v}{B^*(v)} \left(\mathbb{L}\|x_{t_2} - x_{t_1}\| + \mathbb{N}t\|x_{t_2} - x_{t_1}\| + \mathbb{C}T\|x_{t_2} - x_{t_1}\| \right) \\
 & \quad + \frac{v}{B^*(v)\Gamma(v)} \left(\mathbb{L}\|x_{t_2} - x_{t_1}\| + \mathbb{N}t\|x_{t_2} - x_{t_1}\| \right. \\
 & \quad \left. + \mathbb{C}T\|x_{t_2} - x_{t_1}\| \right) \left[\frac{(t_2^v - t_1^v)}{v} \right] + \mu(t)\|x_{t_2} - x_{t_1}\| \\
 & \leq \mu(t)\|x_{t_2} - x_{t_1}\| + \left[\frac{1-v}{B^*(v)} + \frac{(t_2^v - t_1^v)}{B^*(v)\Gamma(v)} \right] \|q(t_2 - t_1)\| \rightarrow 0 \text{ as } t_2 \rightarrow t_1.
 \end{aligned}$$

Similarly, for $t \in (t_k, t_{k+1}]$, $\|\Omega x(t_{k+1}) - \Omega x(t_k)\| \rightarrow 0$ as $t_{k+1} \rightarrow t_k$ where $k = 1, 2, \dots, m$.

With the help of (A4) and (A5), we have proved Ω is equicontinuous.

As result of part (1)-(3) Ω is completely continuous.

Part 4: A priori bounds.

Now it is left to show that that the set

$$\Delta = \{t \in PC^1([- \tau^*, T], \mathbb{R}^n); x = \lambda \Omega(x), \text{ for some } 0 < \lambda < 1\} \text{ is bounded.}$$

Let $x \in \Delta$, for some $0 < \lambda < 1$, for each $t \in [0, T]$, we have $x(t) = \lambda \Omega x(t)$,

$$\begin{aligned}
 x(t) = \lambda & \left[\zeta(0) + \mathcal{F}^*(0) + \sigma(t, x_t) + \frac{1-v}{B^*(v)} \mathcal{B}(t, x_t, Px(t), Qx(t)) \right. \\
 & \left. + \frac{v}{B^*(v)} I_0^v \mathcal{B}(t, x_t, Px(t), Qx(t)) + \sum_{k=1}^m I_k^*(x(t_k^-)) \right].
 \end{aligned}$$

This implies by (A4) and for each $t \in [0, T]$, we have

$$\begin{aligned}
 |x(t)| & = |\lambda \Omega x(t)| \\
 & \leq \lambda |\zeta(0) + \mathcal{F}^*(0)| + \lambda |\sigma(t, x_t)| + \frac{\lambda(1-v)}{B^*(v)} \left| \mathcal{B}(t, x_t, Px(t), Qx(t)) \right| \\
 & \quad + \frac{\lambda v}{B^*(v)\Gamma(v)} \left| \mathcal{B}(t, x_t, Px(t), Qx(t)) \right| I_0^v(t) + \lambda \sum_{k=1}^m \left| I_k^*(x(t_k^-)) \right| \\
 & \leq \lambda |\zeta(0) + \mathcal{F}^*(0)| + \lambda |\sigma(t, x_t)| + \left[\frac{\lambda(1-v)}{B^*(v)} + \frac{\lambda t^v}{B^*(v)\Gamma(v)} \right] \\
 & \quad (\times) \left(\mathbb{L}|x| + \mathbb{N}t|x| + \mathbb{C}T|x| \right) + \lambda \sum_{k=1}^m \left| I_k^*(x(t_k^-)) \right|.
 \end{aligned}$$

Thus by using the inequality (4.1), for every $t \in [0, T]$ and for $\lambda < 1$,

$$\begin{aligned} \|x\| &\leq \lambda\|\zeta(0) + \mathcal{F}^*(0)\| + \lambda\|\sigma(t, x_t)\| + \left[\frac{\lambda(1-v)}{B^*(v)} + \frac{\lambda t^v}{B^*(v)\Gamma(v)} \right] p\|x\| + \lambda m\mathcal{N}^* \\ &\leq \varrho^*. \end{aligned}$$

As it is true for every $x \in \Delta$, Δ is bounded. By Schaefer's fixed point theorem, we claim that Ω has a fixed point which is a solution of the ABC- fractional order impulsive neutral mixed integrodifferential equation (1.3). \square

Theorem 4.3. *If the assumptions (A1) – (A5) hold, then the ABC- fractional order impulsive neutral mixed integrodifferential equation (1.1) has a unique solution, provided that*

$$m\vartheta^* + \mu(t) + \left[\frac{1-v}{B^*(v)} + \frac{t^v}{B^*(v)\Gamma(v)} \right] p < 1 \quad (4.2)$$

where $\sum_{k=1}^m \left\| I_k^*(x_1(t_k)) - I_k^*(x_2(t_k)) \right\| = m\vartheta^*(t)\|x_1 - x_2\|$.

Proof. :We assume the contradiction to prove uniqueness, let x_1 and x_2 be two solutions of (1.1).

Case 1: For $t \in [-\tau^*, 0]$, $\|\Omega x_1 - \Omega x_2\| \rightarrow 0$ implies $x_1 = x_2$.

Case 2: For $t \in [0, t_1]$,

$$\begin{aligned} \|\Omega x_1 - \Omega x_2\| &\leq \mu(t)\|x_1 - x_2\| + \left[\frac{1-v}{B^*(v)} + \frac{t^v}{B^*(v)\Gamma(v)} \right] \\ &\quad (\times) \left(\mathbb{L}\|x_1 - x_2\| + \mathbb{N}t\|x_1 - x_2\| + \mathbb{C}T\|x_1 - x_2\| \right) \\ &\leq \mu(t)\|x_1 - x_2\| + \left[\frac{1-v}{B^*(v)} + \frac{t^v}{B^*(v)\Gamma(v)} \right] p\|x_1 - x_2\|. \end{aligned}$$

Case 3: For $t \in [t_k, t_{k+1}]$,

$$\begin{aligned} \|\Omega x_1 - \Omega x_2\| &\leq \left(\mu(t) + m\vartheta^*(t) \right) \|x_1 - x_2\| + \left[\frac{1-v}{B^*(v)} + \frac{t^v}{B^*(v)\Gamma(v)} \right] \left(\mathbb{L}\|x_1 - x_2\| \right. \\ &\quad \left. + \mathbb{N}t\|x_1 - x_2\| + \mathbb{C}T\|x_1 - x_2\| \right) \\ &\leq \left(\mu(t) + m\vartheta^*(t) \right) \|x_1 - x_2\| + \left[\frac{1-v}{B^*(v)} + \frac{t^v}{B^*(v)\Gamma(v)} \right] p\|x_1 - x_2\|. \end{aligned}$$

By the assumptions (A1) to (A5) and the inequality (4.2), Ω is a contraction. Then by Banach fixed - point theorem, the impulsive neutral FIDE (1.1) has a unique solution. \square

5. Illustration

Consider the following problem

$${}^0_{ABC}D^v \left[x(t) - \frac{e^{-t}x(t)}{(9+e^t)(1+x(t))} \right] = \frac{1}{(t+12)} \frac{x(t)}{1+x(t)} + \int_0^t t\sqrt{2s+1}x(s)ds$$

$$+ \int_0^t s^3 \cos(s^4 - t)x(s)ds$$

$$x(t) = 1 + \phi_0, \quad t \in [-\tau^*, 0], \quad \phi_0 = \sum_{\lambda_i=1}^n \lambda_i x_i$$

$$\Delta x(t_i) = \frac{1}{2}x\left(\frac{1^-}{2}\right), \quad \text{where } t_i = \frac{1}{2}, \quad B^*(v) = 1.$$

By assuming

$$\sigma(t, x_t) = \frac{e^{-t}x(t)}{(9+e^t)(1+x(t))}; Px(t) = \int_0^t t\sqrt{2s+1}x(s)ds;$$

$$Qx(t) = \int_0^t s^3 \cos(s^4 - t)x(s)ds;$$

$$\mathcal{B}(t, x_t, Px(t), Qx(t)) = \frac{1}{(t+12)} \frac{x(t)}{1+x(t)} + \int_0^t t\sqrt{2s+1}x(s)ds$$

$$+ \int_0^t s^3 \cos(s^4 - t)x(s)ds.$$

$$\text{where } \sigma(0, x_0) = \frac{e^{-0}x(0)}{(9+e^0)(1+x(0))} = \frac{0}{(9+1)(1+0)} = \frac{0}{10} = 0;$$

$$\mathcal{B}(0, x_0, Px(0), Qx(0)) = \frac{1}{(0+12)} \frac{x(0)}{1+x(0)} + \int 0\sqrt{2s+1}x(s)ds$$

$$+ \int s^3 \cos(s^4 - 0)x(s)ds.$$

$$\mathcal{B}(0, x_0, Px(0), Qx(0)) = \frac{0}{12} + 0 + \frac{1}{4} \sin 0 = 0$$

Then $\|Px(t) - Px_1(t)\| \leq \frac{1}{3}\|t - t_1\|$ and $\|Qx(t) - Qx_1(t)\| \leq \frac{1}{4}\|t - t_1\|$.

$$\left\| \mathcal{B}(t, x_t, Px(t), Qx(t)) - \mathcal{B}(t_1, x_{t_1}, Px_1(t), Qx_1(t)) \right\|$$

$$\leq \frac{1}{12}\|x(t) - x_1(t)\| + \frac{1}{3}\|x(t) - x_1(t)\|$$

$$+ \frac{1}{4}\|x(t) - x_1(t)\|$$

$$\leq \frac{2}{3}\|x - x_1\|.$$

Hence the conditions A1 – A4 hold true with $\mathcal{L}_1 = \frac{2}{3}$, $\mathbb{N} = \frac{1}{3}$, $\mathbb{C} = \frac{1}{4}$, $\mathcal{N}^* = \frac{1}{2}$,

$\|\sigma(t, x_t)\| \leq \frac{1}{10}$ and $\varrho^* < 1$. Therefore, the above problem has following solution

$$x_n(t) = 1 + \phi_0 + \sigma_{n-1}(t) + (1 - v)f_{n-1}(t) + \frac{1 - v}{\Gamma v} \int_0^t (t - s)^{v-1} f_{n-1}(s) ds.$$

6. Conclusion

The theory of fractional operators with non-singularity is contemporary, and it is essential to study qualitative attributes of mixed integrodifferential equations involving ABC- derivatives. In this paper, we have proved EUS of ABC - fractional dynamical system using Schaefer's and Banach's fixed - point theorem. As a future work, it would be interesting to find new properties and numerical methods of this ABC - fractional operator and some real applications for main results of this paper.

References

1. Abdeljawad, T.: Baleanu, D.: On fractional derivatives with exponential kernel and their discrete versions, Rep. Math. Phys. **80(1)** (2017) 11-27.
2. Abdeljawad, T.: Jarad, F.: Baleanu, D.: On the existence and the uniqueness theorem for fractional differential equations with bounded delay with caputo derivatives, Science in China Series, **51(10)** (2008) 1775-1786.
3. Abdeljawad, T.: Baleanu, D.: Jarad, F.: Existence and uniqueness theorem for a class of delay differential equations with left and right Caputo fractional derivatives, J. Math. Phys. **49(8)** (2008) 083507.
4. Abdeljawad, T.: Baleanu, D.: Integration by parts and its applications of a new non-local fractional derivative with Mittag-Leffler non-singular kernel, J. Nonlinear Sci . **10** (2017) 1098-1107.
5. Abro, K.A.: Khan, I.: MHD flow of fractional Newtonian fluid embedded in a porous medium via Atangana-Baleanu fractional derivatives, Discrete Contin. Dyn. Syst.- S **13(3)** (2020) 377-387.
6. Abro, K.A.: Khan, I.: Nisar, K.S.: Novel technique of Atangana and Baleanu for heat dissipation in transmission line of electrical circuit, Chaos, Solitons & Fractals, **129** (2019) 40-45.
7. Abro, K.A.: Khan, I.: Nisar, K.S.: Use of Atangana-Baleanu fractional derivative in helical flow of a circular pipe, Fractals, **28(8)** (2020) 1-12.
8. Agarwal, R.P.: Zhou, Y.: He, Y.: Existence of fractional neutral functional differential equations, Comput. Math. with Appl. **59** (2010) 1095-1100.
9. Anil Kumar: Muslim Malik: Kottakkaran Sooppy Nisar: Existence and total controllability results of fuzzy delay differential equation with non-instantaneous impulses, Alex. Eng. J. **60(6)** (2021) 6001-6012.
10. Alkhatani, B.S.T.: Chua's Circuit Model with AB derivative with Fractional Order, Chaos, Solitons & Fractals, **89** (2016) 547-551.
11. Al-Refai, M.: Pal, K.: New Aspects of Caputo-Fabrizio Fractional Derivative, Prog. Fract. Differ. Appl. **5(2)** (2019) 157-166.
12. Atangana, A.: Baleanu, D.: New fractional derivative with non-local and non-singular kernel: Theory and application to heat transfer model, Therm. Sci. **20(2)** (2016) 763-769.
13. Atangana, A.: Koca, I.: Chaos in a simple non-linear system with Atangana-Baleanu derivatives with fractional order, Chaos, Solitons & Fractals. **89** (2016) 447- 454.
14. Bainov, D.D.: Simeonov, P.S.: Impulsive Differential Equations: Periodic Solutions and Applications, New York: Longman Scientific and Technical Group. Limited, 1993.
15. Benchohra, M.: Henderson, J.: Ntouyas, S.: Impulsive differential equations and inclusions. New York: Hindawi Publishing Corporation, 2006.
16. Benchohra, M.: Slimani, B.A.: Existence and uniqueness of solutions to impulsive fractional differential equations, Electron. J. Differ. Equ. **2009(10)** (2009) 1-11.

17. Caputo, M.: Fabrizio, M.: A new definition of fractional derivative without singular kernel, *Prog. Fract. Differ. Appl.* **1(2)** (2015) 73-85.
18. Chadha, A.: Pandey, D.N.: Existence results for an Impulsive neutral fractional integro-differential equation with infinite delay, *Int. J. Differ. Equ.* **2014** (2014) ID 780636.
19. Dabas, J.: Chauhan, A.: Existence and uniqueness of mild solution for an impulsive neutral fractional integro-differential equation with infinite delay, *Math. comput. model.* **57(3-4)** (2013) 754-763.
20. Devipriya, R.: Selvi, S.: Solution for ABC-fractional order neutral impulsive differential equation with finite delay, *AIP Conf. Proc.* **2364** (2021) 020005.
21. Esuabana, I.M.: Abasiokwere, U.A.: Formulation of Impulsive Differential Equations with Time -Dependent Continuous Delay, *Am. J. Appl. Math.* **6(4)** (2018) 134-140.
22. Fracisco, J.: Gomez-Aguilar, Atangana, A.: Morales-Delgado, V.F.: Electrical circuits RC, LC and RL described by AB Fractional derivatives, *Int. J. Circuit Theory Appl.* **45(11)** (2017) 1514-1533.
23. Gao, W.: Ghanbari, B.: Bakonous, H.M.: New numerical simulations for some real world problems with Atangana-Baleanu fractional derivative, *Chaos, Solitons & Fractals*, **128** (2019) 34-43.
24. Ghanbari, B.: Atangana, A.: A new application of fractional Atangana-Baleanu derivatives: Designing ABC-fractional masks in image processing, *Phys. A: Stat. Mech. Appl.* **542** (2020) 123516.
25. Guo, T.L.: Jiang, W.: Impulsive fractional functional differential equations, *Comput. Math. with Appl.* **64(10)** (2012) 3414-3424.
26. Jarad, F.: Abdeljawad, T.: Hammouch, Z.: On a class of ordinary differential equations in the frame of Atangana -Baleanu fractional derivative, *Chaos, Solitons & Fractals*, **117** (2018) 16-20.
27. Kaliraj, K.: Thilakraj, E.: Ravichandran, C. : Nisar, K. S.: Controllability analysis for impulsive integro-differential equation via Atangana Baleanu fractional derivative, *Math. Methods Appl. Sci.* (2021) mma.7693.
28. Khan, H.: Jarad, F.: Shah, A.: Khan, A.: Existence and data dependence theorems for solutions of an ABC-fractional order impulsive system, *Chaos, Solitons & Fractals*, **131** (2020) 109477.
29. Khan, H.: Jarad, F.: Abdeljawad, T.: Gomez-Aguilar, A. Khan, Existence and Hyers-Ulam stability for a nonlinear singular fractional differential equations with Mittag-Leffler kernel, *Chaos, Solitons & Fractals*, **127** (2019) 422-427.
30. Khan, H.: Jarad, F.: Abdeljawad, T.: Khan, A.: A singular ABC-fractional differential equation with p-Laplacian operator, *Chaos, Solitons & Fractals*, **129** (2019) 56-61.
31. Koca, I.: Numerical analysis of coupled fractional differential equations with Atangana-Baleanu fractional derivatives, *Discrete Contin. Dyn. Syst.- S* **12(3)** (2019) 475-486.
32. Lakshmikantham, V.: Bainov, D.D.: Simeonov, P.S.: *Theory of impulsive differential equations*. Singapore: World Scientific, 1989.
33. Logeswari, K.: Ravichandran, C.: A new exploration on existence of fractional neutral integro-differential equations in the concept of Atangana- Baleanu derivative, *Phys. A: Stat. Mech. Appl.* **544** (2020) 1-10.
34. Logeswari, K.: Ravichandran, C.: Nisar, K.S.: Mathematical model for spreading of COVID-19 virus with the Mittag Leffler kernel, *Numer. Methods Partial Differ. Equ.* (2020) 22652.
35. Losada, J.: Nieto, J.J.: Properties of a new fractional derivative without singular kernel, *Prog. Fract. Differ. Appl.* **1(2)** (2015) 87-92.
36. Mophou, G.M.: Existence and uniqueness of mild solutions to impulsive fractional differential equations, *Nonlinear Anal-Theor.* **72(3-4)** (2010) 1604-1615.
37. Nuno, R.: Bastos, O.: Calculus of variations involving Caputo-Fabrizio fractional differentiation, *Stat. Optim. Inf. Comput.* **6(1)** (2018) 12-21.
38. Panda, S.K.: Abdeljawad, T.: Ravichandran, C.: Novel fixed point approach to Atangana-Baleanu fractional and L^p -Fredholm integral equations, *Alex. Eng. J.* **59(4)** (2020) 1959-1970.

39. Panda, S.K.: Abdeljawad, T.: Ravichandran, C. A complex valued approach to the solutions of Riemann-Liouville integral, Atangana-Baleanu integral operator and non-linear Telegraph equation via fixed point method, *Chaos, Solitons & Fractals*, **130** (2020) 109439.
40. Panda, S.K.: Ravichandran, C.: Hazarika, B.: Results on system of Atangana Baleanu fractional order Willis aneurysm and nonlinear singularly perturbed boundary value problems, *Chaos, Solitons & Fractals*, **142** (2021) 110390.
41. Podlubny, I.: *Fractional Differential Equations: An Introduction to Fractional Derivatives, Fractional Differential Equations, to Methods of Their Solution and Some of Their Applications*. San Diego: Academic Press, 1999.
42. Ravichandran, C.: Logeswari, K.: Jarad, F.: New results on existence in the frame work of Atangana-Baleanu derivative for fractional integro-differential equations, *Chaos, Solitons & Fractals*, **125** (2019) 194-200.
43. Ravichandran, C.: Logeswari, K.: Panda, S.K.: Nisar, K.S.: On new approach of fractional derivative by Mittag-Leffler kernel to neutral integro-differential systems with impulsive conditions, *Chaos, Solitons & Fractals*, **139** (2020) 110012.
44. Samko, S. G.: Kilbas, A. A.: Marichev, O.I.: *Fractional integrals and derivatives: theory and applications*. Switzerland: Gordon and Breach Science Publishers, 1992.
45. Samoilenko, A. M.: Perestyuk, N. A.: *Impulsive Differential Equations*. Singapore: World Scientific, 1995.
46. Shah, K.: Jarad, F.: Abdeljawad, T.: Mahariq, I.: Qualitative Analysis of a mathematical model in the time of Covid-19, *Biomed - Research International*. **2020** Article ID:5098598 (2020) 11pages.
47. Shaikh, A.: Tassaddiq, A.: Nisar, K.S.: Baleanu, D.: Analysis of differential equations involving Caputo-Fabrizio fractional operator and its applications to reaction-diffusion equations, *Adv. Differ. Equ.* **1** (2019) 1-13.
48. Sheikh, N. A.: Ali, F.: Saqib, M.: Khan, I.: Jan, S. A. A.: Alghamdi, M. S.: Alshomrani, A.S.: Comparison and Analysis of the Atangana-Baleanu and Caputo Fabrizio Fractional derivatives for generalized Casson fluid model with heat generation and chemical reaction, *Results Phys.* **7** (2017) 789-800.
49. Smart, D.R.: *Fixed point theorems*, London: Cambridge University Press, 1980.
50. Tassaddiq, A.: Khan, I.: Nisar, K. S.: Heat transfer analysis in sodium alginate based nanofluid using MoS2 nanoparticles: Atangana-Baleanu fractional model, *Chaos, Solitons & Fractals*, **130** (2020) 109445.
51. Ullah, S.: Khan, M. A.: Farooq, M.: Modelling and analysis of the fractional HBV model with Atangana-Baleanu derivative, *Eur. Phys. J. Plus* **313(12)** (2018) 1-18.
52. Vipin Kumar: Muslim Malik: Controllability results of fractional integro-differential equation with non-instantaneous impulses on time scales, *IMA J. Math. Control.* **38(1)** (2021) 211-231.
53. Vipin Kumar: Muslim Malik: Total controllability and observability for dynamic systems with non-instantaneous impulses on time scales, *Asian J. Control* **23(2)**(2021), 847-859.
54. Zhang, J.: Yin, L.: Zhou, C.: Fractional Herglotz Variational problems with Atangana - Baleanu Fractional Derivatives, *J. Inequal. Appl.* **2018** (2018) 44.
55. Yuldasheva, T.K.: On a Volterra type fractional integro-differential equation with degenerate kernel, *AIP Conf. Proc.* **2365** (2021) 020016.

SOLUTIONS TO ABC-FIDE WITH IMPULSE

R. DEVIPRIYA: PART TIME RESEARCH SCHOLAR, DEPARTMENT OF MATHEMATICS, PERIYAR UNIVERSITY, SALEM - 636 011, INDIA
Email address: devi.jpn@gmail.com

S. SELVI: DEPARTMENT OF MATHEMATICS, N.K.R. GOVERNMENT ARTS COLLEGE FOR WOMEN, NAMAKKAL- 637 001, TAMIL NADU, INDIA
Email address: sselvimaths@yahoo.com

KOTTAKKARAN SOOPPY NISAR: CORRESPONDING AUTHOR: DEPARTMENT OF MATHEMATICS, COLLEGE OF ARTS AND SCIENCES, PRINCE SATTAM BIN ABDULAZIZ UNIVERSITY, WADI AL-DAWASER 11991, SAUDI ARABIA
Email address: n.sooppy@psau.edu.sa, ksnisar1@gmail.com

C. RAVICHANDRAN: DEPARTMENT OF MATHEMATICS, KONGUNADU ARTS AND SCIENCE COLLEGE, COIMBATORE - 641 029, TAMIL NADU, INDIA
Email address: ravibirthday@gmail.com, ravichandran@kongunaducollege.ac.in